

Blinding and Its Quality in Clinical Trials Conducted on Patients with Breast Cancer: A Systematic Review

Abstract

Background: Blinding is one of the critical criteria of clinical trials that prevents probable bias. Judgment regarding results of an intervention significantly depends on the quality of such studies, one of which is blinding. This study aimed to investigate blinding and its quality in clinical trials in patients with breast cancer. **Materials and Methods:** A systematic review was conducted on the online databases of PubMed, ScienceDirect and ProQuest using keywords, MeSH terms and grey literature. Articles were screened by predefined inclusion and exclusion criteria. They were evaluated based on the checklists introduced by Cochrane database. **Results:** From 22519 articles obtained at the initial stage, 20 articles remained after screening for the inclusion and exclusion criteria. Fifteen articles had used single, five: double and none had used triple or quadruple blinding. Seventeen studies had described the details of blinding. Of the 15 single blind articles, the blinded subjects were patients in five, patients and research assistants in three, research assistants in five studies, and two had not given any details. **Conclusions:** The majority of researchers had used the single blind method, though using double, triple or quadruple blinding increases the trustworthiness of results and increases the quality of clinical trials. The details of blinding should be explained to other researchers and for a better understanding of the method if it is to be repeated. Thereafter, nurses can apply new interventions and earn their patients' trust and help those with breast cancer by relieving them of their disease symptoms and its treatment complications.

Keywords: Breast neoplasms, clinical trial, double-blind method

Introduction

Based on recent statistics released by the World Health Organization (WHO), cancer is the second cause of death in the world after cardiovascular disease.^[1] In Iran, it is the third cause of death.^[2] Different interventional studies have been conducted to evaluate the effects of different interventions aimed at relieving the symptoms of the disease and the side-effects of different therapies such as chemotherapy and radiotherapy to reduce morbidity and mortality rates in cancer patients.^[3,4] Any judgment regarding the results of a research and the effect of the intervention involved significantly depends on the quality of such studies.^[5,6]

Compared to other types of studies in health research, a clinical trial is the most appropriate and valuable method for evaluating the effect of a treatment.^[7] High quality clinical trials should be conducted and reported to achieve this goal, and in

doing so allow the audience to judge the internal consistency of the study.^[8] One of the aspects of a clinical trial that indicates the extent to which it is free from probable bias is using the blinding technique.^[6] After controlling the intervention and random allocation, blinding is the most significant criterion in clinical trials, which is included in many quality study tools like the Jadad scale that allocates two fifths of its score to blinding. The Cochrane checklist too is a standard, reliable, and well-known tool for assessing the quality of systematic review articles.^[9,10]

Blinding is one of the methods of reducing the probability of research bias, which can affect the validity of research results.^[7] It is used in different parts of research, such as, concealing data from the participants, data collector and provider, intervention provider, and even data analyzer. Thus, the biases expected to occur in different parts of an interventional study can be

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Matourypour P, Ghorbani A, Mahmoudi M, Binaei N, Manesh HJ, Nayeri ND, *et al.* Blinding and its quality in clinical trials conducted on patients with breast cancer: A systematic review. *Iran J Nurs Midwifery Res* 2022;27:1-7.

Submitted: 03-May-2020. **Revised:** 07-Jul-2020.

Accepted: 16-May-2021. **Published:** 25-Jan-2022.

Pegah
Matourypour¹,
Azam Ghorbani¹,
Mokhtar
Mahmoudi²,
Niloufar Binaei³,
Hadi Jafary
Manesh⁴, Nahid
Dehghan Nayeri⁵,
Imane Bagheri⁶

¹Medical-Surgical Nursing Department, School of Nursing and Midwifery, Tehran University of Medical Science, Tehran, Iran, ²Clinical Care Research Center, Research Institute for Health Development, Kurdistan university of Medical Science, Sanandaj, Iran, ³Department of Adult Health, Faculty of Nursing and Midwifery, Isfahan University of Medical Sciences, Iran, ⁴Department of Medical-Surgical Nursing, School of Nursing and Midwifery, Tehran, Iran, ⁵Nursing and Midwifery Care Research center, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran, ⁶School of Nursing and Midwifery, Isfahan University of Medical Sciences, Isfahan, Iran

Address for correspondence:

Dr. Nahid Dehghan Nayeri,
Nursing and Midwifery Care
Research Center, School of
Nursing and Midwifery, Tehran
University of Medical Sciences,
Tehran, Iran.
E-mail: Dehghann@tums.ac.ir

Access this article online

Website: www.ijnmrjournal.net

DOI: 10.4103/ijnmr.IJNMR_49_20

Quick Response Code:



avoided.^[11,12] In order to investigate blinding, the standards of blinding and its types should be defined first. Blinding refers to concealing information about the type of treatment provided to a specific group of participants.^[12] Simply referring to the types of blinding used is not enough as it can be confusing to the readers. Thus, the authors need to clearly specify who has been blinded in the research process.^[7] The accurate interpretation of a clinical trial is possible when there is accurate information on the methods of design and analysis of outcomes.^[13] Previous studies indicate that blinding is one of the issues less addressed in clinical trial designs.^[13-16]

Subsequently, as mentioned above, a variety of biases ensue and the study results become questionable.^[7] The interpretation of clinical trials' results depends on the quality of methods and blinding as a means to prevent bias. Given that breast cancer is the most prevalent cancer and second cause of mortality in women,^[3] a search was conducted in the Cochrane database to check whether the subject was not repetitive. We found several studies that had examined the quality of clinical trials but not specifically in patients with breast cancer.^[11,13,15] Hence, in this study we aimed to investigate blinding and its quality in clinical trials in patients with breast cancer.

Materials and Methods

This research is a systematic review conducted between 2012 and 2019. The search strategy, article selection, and evaluation of articles' quality are explained in details below: Search was conducted in the valid medical science databases of PubMed with the MeSH term 'single blind method' and 'double blind study' and, 'breast neoplasm' and other databases such as ScienceDirect and ProQuest with the keywords of breast cancer/neoplasm and blind study and single/double blind study on clinical trials in humans as well as the Iranian database 'SID' with the keywords 'breast cancer' and 'Blinding' [Table 1]. Then, the articles were evaluated based on the checklists introduced by the Cochrane database.^[17]

The inclusion criteria included, the presence of the keywords 'breast cancer' and 'blinding' in the title and abstract, the interventional nature of the research, English or Persian language articles, the intervention being conducted in the nursing field. The exclusion criteria included, not having used blinding methods, the acquisition of less than 12 points from the checklist, and the lack of availability of the articles' full texts.

During the first stage, all the articles were reviewed by three researchers in terms of relevancy of the titles and abstracts (interventional method/application of blinding/patients with breast cancer) and irrelevant articles were removed. Then, the articles' full texts were acquired and after omitting the names of the authors and the journals the reviewers began reviewing the articles.

Thereafter, the section on the type and manner of blinding by the corresponding checklist was completed for each English and Persian article. Obtaining at least 12 scores from 20 was mandatory for inclusion.^[17,18] Finally, an expert on research methodology evaluated the assessment procedures on the final articles and made suggestions to be applied. After the first search we screened the final articles' references as our second search, but no new article was found.

The checklist was designed based on the items introduced by Cochrane for systematic reviews.^[17] Its validity and reliability were measured and confirmed by ten experts in the research methodology, epidemiology, and nursing groups. The calculated Content Validity Index (CVI) was 0.87 and the reliability of all the items on the checklist was estimated at 0.92 Cronbach's Alpha, both of which were acceptable. The checklist is demonstrated in Table 2.

Ethical considerations

This study was approved by the ethics committee of the Nursing and Midwifery Research Centre of Tehran University of Medical Sciences, Tehran, Iran (IR.TUMS.FNM.REC.1399). In this systematic review and meta-analysis, the collected data were only used for scientific purposes, and intellectual property was observed in the reporting and publication of the results.

Results

Of the 22519 articles retrieved at the initial stage of the search, first the duplicate articles were removed (6832 articles). Twenty articles were finally included in the review based on the inclusion and exclusion criteria and checklist scores [Figure 1]. The final results are shown in Table 2 in terms of the intervention, outcome, type of blinding, sample and randomization.

Target population

Clinical trials conducted on various types of patients with breast cancer^[19-38] were included, the risk of breast cancer incidence among this population,^[26,29,32] breast cancer survivors,^[22,23,27,28,33-35] those suffering from breast cancer,^[19,21,24,25,30,31,36-38] patients undergoing chemotherapy,^[24,25,38] patients undergoing radiotherapy,^[21] and patients awaiting surgery.^[20]

Studied variables

In the final stage, the effects of different interventions were evaluated to decrease or eliminate certain variables including fatigue,^[19,22,23,34] musculoskeletal symptoms,^[27,30,36] conditional and acute nausea and vomiting due to chemotherapy (intensity, duration),^[24,25,38] interventions on lifestyle,^[32] physical and mental quality of life, cognitive function and physical activity,^[23] physical function,^[19,27] the incidence of breast cancer,^[29] mental stress,^[27,28] cognitive function (memory and information processing speed),^[35]

Table 1: Search strategy

Search term	PubMed	ScienceDirect	ProQuest	SID*
(Single-blind method or double- blind study) and breast neoplasm	642			30
MeSH				
(Breast cancer OR neoplasm) and blind study and (single and double study)	321	4941	16585	
Keywords				
Total	963	4941	16585	30

*SID: Scientific Information Database




Figure 1: Figure showing the different phases involved in searching relevant publications

preoperative anxiety,^[20] treatment-related symptoms (pain and stress, radiotherapy),^[21] lymphedema,^[31] remembering routine screening times,^[26] cellular immunity,^[36] and depression.^[37]

Interventions in clinical trials

Interventions conducted in different clinical trials included, therapeutic touch (energy therapy),^[24,25,38] short-term psychotherapy,^[37] Chinese medicinal herbs,^[36] physical therapy,^[31] Tai Chi and Qigong Day,^[22,23] SMS reminders,^[26] short dance moves,^[21] aromatherapy,^[20] cognitive therapy,^[35] acupressure,^[34] recognition of behavioral therapy,^[28] support groups,^[33] green tea extract consumption,^[20] acupuncture,^[30] weight lifting,^[27] progressive muscle relaxation and mindfulness meditation^[19] and lifestyle interventions.^[32]

Types of blinding and their quality

From a total of 20 articles, 15 had used single, 5 had used double and neither study had used triple or quadruple

blinding. Of the 15 single blind articles, the blinded subjects were patients in five studies^[20,21,24,25,38]; they were patients and research assistants in three studies^[33-35] and were research assistants in five studies.^[18,19,27,28,32] Two studies failed to explain the blinded subjects and their details.^[26,31] In five articles, double blinding was used.^[22,23,29,30,36] In these studies, except for one study, the details of blinding had not been mentioned.^[36] In the remaining four cases, blinding had been performed on the samples and the researcher^[29] and the samples and subjects, who analyzed data^[22,23] on the samples and the physician/nurse/other member of the treatment team^[30] were unaware of the intervention type.

Discussion

Of the 20 final articles retrieved from nursing – related studies, 15 had used the single blind method while five studies had used double blinding. No study had used triple or quadruple blinding. This is not a desirable finding, as unintentional systematic bias may occur and it can threaten the reliability of the research results, which can be minimized by blinding. Polit and Beck introduced blinding as a technique that can reduce bias through single, double, triple and quadruple blind methods.^[7] The greater the number of blinded parties in the research, the lower the probability of bias that is out of the researcher’s control. Thus, indicating the importance of employing robust methods in clinical trials to raise the validity of results.

Of the 15 single-blind articles, the blinded subjects were patients in five studies^[20,21,24,25,38]; they were patients and research assistants in three studies^[33-35] and were research assistants in five studies.^[18,19,27,28,32] Two studies failed to give any explanations regarding the blinded subjects and their details.^[26,31] In five articles, the double blind method was used.^[22,23,29,30,36] Seventeen out of 20 studies had described the details of their blinding, although some studies had not done so. The quality with which clinical trials are conducted and describing blinding along with its details are of great importance, as nurses can use reliable research results that can improve patient care. Therefore, blinding along with other positive advantages of the method like multi-group random allocation, and allocation concealment can increase the quality of a clinical trial and yield more accurate and reliable results. Paying attention to blinding improves the quality of such

Table 2: Clinical trials conducted on breast cancer in the field of nursing that have used the blind method

Authors	Year	Type of Blind Method	Intervention	Outcome (s) Assessed
Gok Z <i>et al.</i> ^[19]	2019	Single The assessor was blinded	The participants were randomly assigned to either a 12-week PMR* or MM** intervention or to the Control Group (CG). The intervention group continued PMR or MM for 20 minutes every day, for a total of 12 weeks. The CG received only a single attention-matched educational session (15 min) on breast cancer before the start of the paclitaxel regimen. Data were collected at baseline, week 12, and week 14.	Fatigue, Functional Living Index-Cancer (FLIC)
Franco L <i>et al.</i> ^[20]	2016	Single Only the patients were blinded	In the preoperative holding area, subjects received 2 drops of oil, either 2% LFO (lavender fleur oil) or UO (unscented oil), inside a plastic oxygen face mask for 10 min	Anxiety and vital signs before and after aromatherapy
Ho RTH <i>et al.</i> ^[21]	2016	Single Only the participants were blinded to the group allocation.	The intervention included six 1.5-h DMT*** sessions provided twice a week over the course of radiotherapy.	Perceived stress, anxiety, depression, fatigue, pain, sleep disturbance, and quality of life
Larkey L <i>et al.</i> ^[22]	2016	Double The participants and study staff involved in data collection and/or analysis were blinded to the assignment.	The intervention group (QG/TCE****, SQG*****, ES*****) met weekly for an hour during the 12 weeks of the intervention.	The primary outcome (fatigue) and secondary outcomes (anxiety, depression, sleep quality, cognitive function, and physical activity) were assessed at baseline, immediately and 6 months after the intervention.
Larkey LK <i>et al.</i> ^[23]	2016	Double The participants and study staff involved in data collection and/or analysis were blinded.	Twelve weekly sessions of QG/TCE were compared to sham Qigong (SQG), a gentle movement control intervention similar to QG/TCE but without focusing on breathing and the meditative state.	Mental and physical QOL, cognitive function (Functional Assessment of Cancer Therapy-Cognitive Function, overall levels of physical activity and body mass index (BMI) were assessed at 3 time points.
Matourypour <i>et al.</i> ^[24]	2016	Single Patients were not aware of the intervention (placebo group).	Therapeutic touch was applied to each patient once for 20 min on the aura (human energy field) focusing on the solar chakra.	Chemotherapy-induced vomiting
Vanaki <i>et al.</i> ^[25]	2016	Single Patients were not aware of the intervention (placebo group).	Therapeutic touch was carried out for both (test and placebo) groups prior to their first chemotherapy appointment (once for about 15-20 minutes).	Nausea duration and frequency
Kerrison <i>et al.</i> ^[26]	2015	Single	Delivery of a text-message reminder 48 h before appointment, which included the time, date and venue of the appointment, as well as information about rescheduling if unable to attend.	Breast screening end codes at the initial appointment and again 60 days thereafter.
Brown JC <i>et al.</i> ^[27]	2015	Single Measurements were obtained by trained staff who were blinded to the study group assignments.	Twice-weekly slowly progressive weight lifting or standard care.	Physical function
Stagl <i>et al.</i> ^[28]	2015	Single Randomization and assessment were conducted by blinded study coordinators.	A 10-week, group-based Cognitive Behavioral Stress Management (CBSM) intervention (<i>n</i> =120) for the test group and/or a 1-day psychoeducational seminar for the control group (<i>n</i> =120).	Survival and recurrence 8-15 years post-enrollment.

Contd...

Table 2: Contd...

Authors	Year	Type of Blind Method	Intervention	Outcome(s) Assessed
Dostal AM <i>et al.</i> ^[29]	2015	Double Participants, investigators, laboratory staff, and those monitoring clinical outcomes and adverse events were blinded to treatment assignments.	Administration of four oral GTE, ^{††} capsules containing 1315 mg±116 total catechins per day (843±44 mg as Epigallocatechin-3-gallate (EGCG)) for 12 months.	Mammographic density, circulating reproductive hormones, and biomarkers of breast cancer risk.
Bao <i>et al.</i> ^[30]	2014	Double The treating oncologist (s), nurses, and study team members. The participants too were blinded to their treatment assignments.	Patients were randomized to an 8-week-long Real Acupuncture (RA) group or Sham Acupuncture (SA) group.	Menopausal symptoms, depression, anxiety and depression, sleep quality, quality-of-life.
Tambour M <i>et al.</i> ^[31]	2014	Single	Group A: Complete Decongestive Therapy including manual drainage Group B: Complete Decongestive Therapy without manual lymphatic drainage.	Lymphedema (%) from baseline to 7 months, body weight (kg), patient sensation of heaviness, sensation of tension and quality of life.
Anderson <i>et al.</i> ^[32]	2014	Single The researcher was blinded to the participant's group allocation.	Intervention participants were scheduled for a one-hour lifestyle coaching session (face-to-face) and up to six fortnightly follow-up telephone consultations for three months	Body weight, waist circumference, eating and alcohol habits, and physical activity
Carlson <i>et al.</i> ^[33]	2013	Single Participants as well as research assistant were blinded to the condition at the baseline assessment.	Mindfulness-Based Cancer Recovery (MBCR) consisted of eight weekly group sessions, 90 minutes each, plus a 6-hour workshop between weeks 6 and 7 for a total of 18 contact hours. Supportive expressive therapy consisted of 12 weekly group sessions, 90 minutes each, and equal contact hours for MBCR.	Mood, stress symptoms, quality of life, social support, spirituality and post-traumatic growth immediately before and after the interventions, and 6 and 12 months later.
Zick <i>et al.</i> ^[34]	2012	Single Participants, investigators, study nurses and all study staff were blinded.	6-weeks of relaxation acupressure compared to stimulatory acupressure or standard care.	Fatigue, depression, anxiety, self-efficacy, sleep quality.
Von Ah <i>et al.</i> ^[35]	2012	Single The participants and cognitive testers were blinded	Working memory and processing speed educational sessions, including ten 1-hour training sessions conducted in small groups of 3-5 breast cancer survivors over 6-8 weeks.	Working memory and processing speed, perceived cognitive functioning, anxiety, fatigue, quality of life, intervention satisfaction, acceptability.
Zhuang SR <i>et al.</i> ^[36]	2012	Double	Chinese medicinal herbs	Cellular immunity.
Zwerenz <i>et al.</i> ^[37]	2012	Single Assessments were done by independent, trained and supervised research-assistants, who were blinded to the intervention.	Short-term psychodynamic psychotherapy (up to 20±5 sessions).	Depression and quality of life
Matoury <i>et al.</i> ^[38]	2013	Single Patients were not aware of the intervention (placebo group).	Therapeutic touch program on women with breast cancer under chemotherapy.	Nausea

*PMR: Progressive Muscle Relaxation, **MM: Mindfulness Meditation, ***DMT: Dance Movement Therapy, ****QG/TCE: Qigong/Tai Chi Easy, *****ES: educational support, *****SQM: "sham" Qigong group (movements without a focus on the breath and meditative state), ^{††}GTE: Green Tea Extract

studies because of their significant impact on patient care. In fact, attention is paid to the quality of studies in terms of design to enable the generalization of their results, at the same time that analyzing the quality of the reported

findings is of great value to the readers and/or users of research results.^[15]

Since blinding is one of the criteria for evaluating the quality of research, double or even triple or quadruple blind methods are recommended. Given that this kind of methodology has higher quality the results of a clinical trial that has employed this methodology can be trusted more too. Thus, future overviews on other factors affecting the quality of the methods employed, such as sampling, sample size calculation formula, data collection, etc., are also suggested.

This research has certain limitations. We did not have access to the full texts of some of the articles, despite the correctness of their titles and abstracts and their suitability for inclusion in the review. Moreover, limiting the language of the articles to English/Persian and restricting our search to four databases were other limitations of our study, whereas, other databases could contain further relevant and valuable researches.

Conclusion

Blinding has a critical role in preventing bias, so using double, triple or quadruple blinding increases the trustworthiness of clinical trial results. Thus, in any clinical trial, it should be clearly specified who has been blinded to raise the trustworthiness of results. Thereafter, nurses can apply new interventions and earn their patients' trust and help those with breast cancer by relieving them of their disease symptoms and its treatment complications by using such trustworthy results.

Acknowledgments

This research has been approved by the Nursing and Midwifery Care Research Center of Tehran University of Medical sciences under project number 47816.

Financial support and sponsorship

Tehran University of Medical Sciences

Conflicts of interest

Nothing to declare.

References

- World Health Organization. Noncommunicable Disease, countries profile. 2018. Available from: <https://www.who.int/nmh/publications/ncd-profiles-2018/en/>. [Last access on 2021 Aug 24].
- Matourypour P. Cancer from Nursing Perspective. Tehran: Heidari Publication; 2017.
- Smeltzer SC, Bare BG, Hinkle JL, Cheere KH, Kluwer W. Brunner and Sudarth's Textbook of Medical-Surgical Nursing. 16th ed, Philadelphia. Lippincot Williams and Wilkins; 2018.
- Matourypour P. Clinical Guideline for Nursing Students and Nurses in Patients with Breast Cancer. Tehran: Heidari Publication; 2016.
- Shahsavari H, Matory P, Zare Z, Taleghani F, Kaji MA. Effect of self-care education on the quality of life in patients with breast cancer. *J Edu Health Promot* 2015;4:70.
- Burns N, Grove S. The practice of Nursing Research. 9th ed, Riverport Lane. Elsevier; 2020.
- Polit DF, Beck CT. Nursing research: Generating and assessing evidence for nursing practice. 10th ed. Philadelphia: Williams and Wilkins; 2017.
- Mehrazmay A, Karambakhsh A, Salehi M. Reporting quality assessment of randomized controlled trials published in nephrology urology monthly journal. *Nephrourol Mon* 2015;7:e28752.
- Alam M, Rauf M, Ali S, Nodzinski M, Minkis K. A systematic review of reporting in randomized controlled trials in Dermatologic Surgery: Jadad scores, power analysis, and sample size determination. *Dermatol Surg* 2014;40:1299-305.
- McCormick F, Cvetanovich GL, Kim JM, Harris JD, Gupta AK, Abrams GD, *et al.* An assessment of the quality of rotator cuff randomized controlled trials: Utilizing the Jadad score and CONSORT criteria. *J Shoulder Elbow Surg* 2013;22:1180-5.
- Khatami A, Firooz A. Evaluation of clinical trials. *Iran J Dermatol* 2008;11:76-86.
- Polit DF, Beck CT. Nursing research: Generating and assessing evidence for nursing practice. 9th ed. Philadelphia: Williams and Wilkins; 2013.
- Ayatollahi MT, Jafari P, Ghaem H. Evaluation of the quality of clinical trials published in Iranian medical journals between the years 2001-2002. *J Babol Univ Med Sci* 2004;7:64-70.
- Kenneth MD, Schulz KF, Altman DG. The CONSORT statement: Revised recommendations for improving the quality of reports of parallel-group randomised trials. *Clin Oral Investig* 2003;7:2-7.
- Saurav G, Eunjung K, Wonku K, Eunyung K. Assessment of adherence to the CONSORT statement for quality of reports on randomized controlled trial abstracts from four high-impact general medical journals. *Trials* 2012;13:77.
- Chan A-W, Altman DG. Identifying outcome reporting bias in randomised trials on PubMed: Review of publications and survey of authors. *BMJ* 2005;330:753.
- Dostii M, Shahsavari H, Zare Z, Talghani F. Systematic Reviews of Studies on Effectiveness: Help Center Review and Dissemination (CRD) for Performers, Isfahan. Isfahan University of Medical Sciences and Health Services publication; 2010
- Matory P, Gholami R, Dehghan M, Vanaki Z, Shirazi M, Binaee N, *et al.* Efficacy of complementary therapies in reduction of chemotherapy induced nausea and vomiting in breast cancer patients: Systematic review. *Complement Med* 2014;4:831-44.
- Gok Metin Z, Karadas C, Izgu N, Ozdemir L, Demirci U. Effects of progressive muscle relaxation and mindfulness meditation on fatigue, coping styles, and quality of life in early breast cancer patients: An assessor blinded, three-arm, randomized controlled trial. *Eur J Oncol Nurs* 2019;42:116-25.
- Franco L, Blanck TJJ, Dugan K, Kline R, Shanmugam G, Galotti A, *et al.* Both lavender fleur oil and unscented oil aromatherapy reduce preoperative anxiety in breast surgery patients: A randomized trial. *J Clin Anesth* 2016;33:243-9.
- Ho RTH, Fong TCT, Cheung IKM, Yip PSF, Luk M-Y. Effects of a short-term dance movement therapy program on symptoms and stress in patients with breast cancer undergoing radiotherapy: A randomized, controlled, single-blind trial. *J Pain Symptom Manage* 2016;51:824-31.
- Larkey L, Huberty J, Pedersen M, Weihs K. Qigong/Tai Chi Easy for fatigue in breast cancer survivors: Rationale and design of a randomized clinical trial. *Contemp Clin Trials* 2016;50:222-8.
- Larkey LK, Roe DJ, Smith L, Millstine D. Exploratory outcome

- assessment of Qigong/Tai Chi Easy on breast cancer survivors. *Complement Ther Med* 2016;29:196-203.
24. Matourypour P, Vanaki Z, Zare Z, Mehrzad V, Dehghan M, Ranjbaran M. Investigating the effect of therapeutic touch on the intensity of acute chemotherapy-induced vomiting in breast cancer women under chemotherapy. *Iran J Nurs Midwifery Res* 2016;21:255-60.
 25. Vanaki Z, Matourypour P, Gholami R, Zare Z, Mehrzad V, Dehghan M. Therapeutic touch for nausea in breast cancer patients receiving chemotherapy: Composing a treatment. *Complement Ther Clin Pract* 2016;22:64-8.
 26. Kerrison RS, Shukla H, Cunningham D, Oyeboode O, Friedman E. Text-message reminders increase uptake of routine breast screening appointments: A randomised controlled trial in a hard-to-reach population. *Br J Cancer* 2015;112:1005-10.
 27. Brown JC, Schmitz KH. Weight lifting and appendicular skeletal muscle mass among breast cancer survivors: A randomized controlled trial. *Breast Cancer Res Treat* 2015;151:385-92.
 28. Stagl JM, Lechner SC, Carver CS, Bouchard LC, Gudenkauf LM, Jutagir DR, *et al.* A randomized controlled trial of cognitive-behavioral stress management in breast cancer: Survival and recurrence at 11-year follow-up. *Breast Cancer Res Treat* 2015;154:319-28.
 29. Dostal AM, Samavat H, Bedell S, Torkelson C, Wang R, Swenson K, *et al.* The safety of green tea extract supplementation in postmenopausal women at risk for breast cancer: Results of the Minnesota Green Tea Trial. *Food Chem Toxicol* 2015;83:26-35.
 30. Bao T, Cai L, Snyder C, Betts K, Tarpinian K, Gould J, *et al.* Patient-reported outcomes in women with breast cancer enrolled in a dual-center, double-blind, randomized controlled trial assessing the effect of acupuncture in reducing aromatase inhibitor-induced musculoskeletal symptoms. *Cancer* 2014;120:381-9.
 31. Tambour M, Tange B, Christensen R, Gram B. Effect of physical therapy on breast cancer related lymphedema: Protocol for a multicenter, randomized, single-blind, equivalence trial. *BMC Cancer* 2014;14:239.
 32. Anderson AS, Macleod M, Mutrie N, Sugden J, Dobson H, Treweek S, *et al.* Breast cancer risk reduction-Is it feasible to initiate a randomised controlled trial of a lifestyle intervention programme (ActWell) within a national breast screening programme? *Int J Behav Nutr Phys Act* 2014;11:156.
 33. Carlson LE, Doll R, Stephen J, Faris P, Tamagawa R, Drysdale E, *et al.* Randomized controlled trial of Mindfulness-based cancer recovery versus supportive expressive group therapy for distressed survivors of breast cancer. *J Clin Oncol* 2013;31:3119-26.
 34. Zick SM, Wyatt GK, Murphy SL, Arnedt JT, Sen A, Harris RE. Acupressure for persistent cancer-related fatigue in breast cancer survivors (AcuCrft): A study protocol for a randomized controlled trial. *BMC Complement Altern Med* 2012;12:132.
 35. Von Ah D, Carpenter JS, Saykin A, Monahan P, Wu J, Yu M, *et al.* Advanced cognitive training for breast cancer survivors: A randomized controlled trial. *Breast Cancer Res Treat* 2012;135:799-809.
 36. Zhuang SR, Chiu HF, Chen SL, Tsai JH, Lee MY, Lee HS, *et al.* Effects of a Chinese medical herbs complex on cellular immunity and toxicity-related conditions of breast cancer patients. *Br J Nutr* 2012;107:712-8.
 37. Zwerenz R, Beutel ME, Imruck BH, Wiltink J, Haselbacher A, Ruckes C, *et al.* Efficacy of psychodynamic short-term psychotherapy for depressed breast cancer patients: Study protocol for a randomized controlled trial. *BMC Cancer* 2012;12:578.
 38. Matoury P, Vanaki Z, Zare Z, Mehrzad V, Dehghan M. Investigation of the effects of therapeutic touch on intensity of anticipatory and acute induced-nausea in breast cancer women undergoing chemotherapy in Isfahan. *Complement Med J* 2013;3:585-94 (Persian).